

International Journal of Physiology, Health and Physical Education



ISSN Print: 2664-7265
ISSN Online: 2664-7273
Impact Factor: RJIF 8
IJPHE 2024; 6(2): 120-123
www.physiologyjournals.com
Received: 08-07-2024
Accepted: 15-08-2024

Dr. Amit S Patel
Incharge Principal, MB Gohil
Institute of Medical Science &
Research Center, College of
Physiotherapy, Navsari,
Gujarat, India

Correlation between obesity and cardiopulmonary capacity in school going children of Navsari city

Dr. Amit S Patel

DOI: <https://doi.org/10.33545/26647265.2024.v6.i2b.80>

Abstract

Background: Childhood obesity is a burgeoning global public health concern, posing significant risks to cardiovascular and respiratory health. This study aimed to explore the correlation between obesity and cardiopulmonary capacity among school-going children aged 8 to 16 years in Navsari, India.

Methods: A correlational study was conducted, involving a convenient sample of 55 school children. Body mass index (BMI) was calculated, and participants underwent the 6-minute walk test (6MWT) to evaluate cardiopulmonary capacity. Maximum oxygen uptake (VO_{2max}) was estimated utilizing a prediction equation based on 6MWT distance and BMI.

Results: Analysis revealed a noteworthy negative correlation between BMI and VO_{2max} ($r = -0.874$, $p < 0.001$), indicating that heightened obesity levels corresponded to diminished cardiorespiratory fitness. The 6MWT emerged as a dependable and valid measure for assessing aerobic capacity in obese youth.

Conclusion: In conclusion, the study on the correlation between obesity and cardiopulmonary capacity in school-going children of age 8 to 16 years of Navsari city, India; reinforced strong negative correlation between BMI and maximum oxygen uptake (VO_{2max}), indicating that higher levels of obesity were associated with reduced cardiopulmonary capacity.

Keywords: Childhood obesity, cardiopulmonary capacity, VO_{2max} , 6-minute walk test, body mass index, school children

Introduction

Childhood obesity has emerged as a pressing global health concern, with its prevalence escalating at an alarming rate in both developed and developing nations. The World Health Organization has identified childhood obesity as one of the most serious public health challenges of the 21st century [1]. In India, the prevalence of obesity among children and adolescents has risen significantly in recent decades, driven by factors such as sedentary lifestyles, unhealthy dietary habits, and rapid urbanization [2].

Obesity in childhood is associated with numerous adverse health consequences, including an increased risk of cardiovascular diseases, type 2 diabetes, and respiratory disorders [3]. Of particular concern is the impact of obesity on cardiopulmonary capacity, which refers to the ability of the cardiovascular and respiratory systems to supply oxygen during physical activity [4]. Reduced cardiopulmonary capacity can lead to decreased exercise tolerance, fatigue, and a diminished quality of life, further exacerbating the negative consequences of obesity [5].

Assessing cardiopulmonary capacity is crucial for identifying individuals at risk and developing appropriate interventions. The six-minute walk test (6MWT) has emerged as a reliable and widely used method for evaluating functional exercise capacity in various populations, including obese children and adolescents [6,7]. The 6MWT measures the distance an individual can walk on a flat surface in six minutes, providing an estimate of their maximal oxygen uptake (VO_{2max}), a key indicator of cardiopulmonary fitness [8].

Despite the increasing prevalence of childhood obesity and its potential impact on cardiopulmonary health, there is a paucity of research investigating the correlation between obesity and cardiopulmonary capacity in specific geographic regions. Understanding this relationship within a local context is essential for developing targeted interventions and preventive strategies tailored to the unique needs and characteristics of the community [9]. The present study aims to address this gap by investigating the correlation between obesity and cardiopulmonary capacity among school-going children aged 8 to 16 years in Navsari, India.

Corresponding Author:
Dr. Amit S Patel
Incharge Principal, MB Gohil
Institute of Medical Science &
Research Center, College of
Physiotherapy, Navsari,
Gujarat, India

Methodology

- **Study design:** Correlational study.
- **Sample design:** Convenient sampling.
- **Sample population:** School-going children aged 8 to 16 years both male and female.
- **Sample size:** 55 [6]. (As per the time-bound nature of the study).
- **Study setting:** Schools located in Navsari city.
- **Study duration:** 6 Months.

Material Used

- Measurement scale in meter.
- Non-adhesive measure tape.
- Weighing machine.
- Two small cones to mark the turnaround points.
- A chair that can be easily moved along the walking course.
- Countdown Timer (or stopwatch).
- Calculator or Mobile phone.
- Chalk.
- Red marking at the starting point.
- Worksheets on a clipboard.
- First aid.
- BMI graph [20].

Sampling Criteria

Inclusion Criteria

- Children aged between 8 to 16 years both male and female [2].
- Body Mass Index (BMI) above the 95th percentile for age and sex, based on the WHO growth reference standards [2].
- Ability to comprehend and adhere to instructions [2].
- Consent signed from parents or guardian or caregivers. [2].
- Willing participants.

Exclusion Criteria

- Children diagnosed with metabolic syndrome, severe/chronic cardiac, or neurological disorders [2].
- History of acute or chronic cardiopulmonary diseases [2].
- Previous history of musculoskeletal injuries [2].
- Children with significant coordination and balance disorders predisposing them to falls, as determined by a comprehensive medical history [2].

- Those already engaged in exercises other than routine exercises [2].

Result

The study involved 55 school-going children aged between 8 to 16 years from Navsari city. Their cardiopulmonary capacity, represented by VO₂max, was assessed using the 6-minute walk test (6MWT).

The data for BMI and VO₂ max were normally distributed, while the data for 6MWD was not normally distributed.

Table 1: Describes variables including Mean, Median, Standard deviation of BMI, VO₂max and Distance

Variables	Mean ± SD
BMI	29.921 ±3.99169
VO ₂ max	22.8438±1.99598
Distance	5.2284±7.01368

The mean BMI of the participants was 29.93 kg/m², which falls in the obese category. The mean VO₂max was 22.84 ml/kg/min.

Table 2: The normality of data distribution including BMI, VO₂max & distance

	Kolmogorov- Smirnov		
	Statistic	Df	Sig.
BMI	0.102	55	0.200*
VO ₂ max	0.65	55	0.200*
Distance	0.127	55	0.027

*. This is a lower bound of the true significance.

Table 2 represents the normality of the data distribution indicate that the data for BMI and VO₂max were normally distributed (*p* > 0.05), while the data for distance was not normally distributed (*p* = 0.027).

Table 3: Relationship between obesity (BMI) and cardiopulmonary capacity (VO₂max), Pearson’s correlation coefficient was calculated

Variable		VO ₂ max
BMI	Pearson Correlation (r)	-.874**
	Sig. (2-tailed) (p Value)	.000
	N	55

***. Correlation is significant at the 0.01 level (2 tailed)

Table 3 reveals There was a strong negative correlation between BMI and VO₂max (*r* = -0.874, *p* < 0.001).




Fig 1: Relationship between obesity (BMI) and cardiopulmonary capacity (VO₂max), were calculated and scatter diagram was plotted accordingly

This indicates that as the children's BMI (obesity level) increased, their VO₂ max (cardiopulmonary capacity) decreased.

Table 4: Examines the relationship between VO₂ max and 6MWD of the participants

Variable		Distance
VO ₂ max	Pearson Correlation (r)	.671**
	Sig. (2-tailed) (p Value)	.000
	N	55

** . Correlation is significant at the 0.01 level (2 tailed)

Table 4 reveals there was a positive correlation between VO₂max and the 6-minute walk distance (6MWD) ($r =$

0.671, $p < 0.001$), as both are indicators of cardiorespiratory fitness.




Fig 2: Represents the relationship between VO₂max and 6MWT of the participants

The above graph represents positive correlation between VO₂max and 6MWD as both measures are indicators of cardiorespiratory fitness.

The data for BMI and VO₂max were normally distributed, while the data for 6MWD was not normally distributed.

Discussion

The results of this study on the correlation between obesity and cardiorespiratory capacity in school-going children of Navsari city are consistent with findings from several other studies in this area.

The study's primary finding was a strong negative correlation between body mass index (BMI) and maximum oxygen uptake (VO₂max) among the obese school children ($r = -0.874$, $p < 0.001$). This inverse relationship between increased adiposity and reduced cardiorespiratory fitness is well-documented in the literature.

Similar findings have been reported by Grzyb *et al.* (2021), who observed that obese children and adolescents exhibited significantly lower VO₂ peak values relative to their body weight when compared to normal-weight and overweight peers [6]. Dobrowolska *et al.* (2022) also found a significant correlation between fat-free mass and absolute aerobic capacity (VO₂ peak) in children and youth with elevated BMI [12].

The diminished cardiorespiratory capacity in obese children is attributed to the increased metabolic demand placed on the cardiovascular and respiratory systems during physical exertion, leading to premature fatigue and reduced exercise tolerance [6, 15]. Additionally, obesity is often accompanied by sedentary lifestyles and decreased levels of physical

activity, further exacerbating the decline in cardiorespiratory fitness [6].

The present study employed the 6MWT as the assessment tool to evaluate cardiorespiratory capacity, aligning with the recommendations in the literature. The 6MWT has been widely used and validated for assessing functional exercise capacity in various populations, including obese children and adolescents [9, 10].

Morinder *et al.* (2008) and Li *et al.* (2005) have previously established the 6MWT as a reliable and valid test for evaluating exercise tolerance and endurance in obese youth and healthy children, respectively [9, 10].

Vanhelst *et al.* (2013) demonstrated that the distance walked during the 6MWT and BMI are the most significant factors in predicting VO₂max in obese youth, highlighting the utility of this prediction equation [11]. Interventions targeting weight management and 34 increased physical activity levels have the potential to prevent or ameliorate the adverse consequences of obesity on cardiorespiratory health and overall well-being in children [6, 14].

As noted by Ahirwar and Mondal (2019), childhood obesity has become a global public health crisis, with a rising prevalence driven by various factors, such as sedentary lifestyles and unhealthy dietary habits [2]. Tackling this issue through comprehensive, community-based strategies is essential to safeguard the long-term cardiovascular and respiratory health of children.

In conclusion, the present study's findings are well-aligned with the existing literature, reinforcing the negative impact of obesity on cardiorespiratory fitness in school-going children. The consistent use of reliable assessment tools, such as the 6MWT, and the application of predictive

equations further strengthen the comparability and validity of the results across these studies. The collective evidence underscores the urgency of addressing childhood obesity through targeted interventions and preventive measures to promote overall health and well-being in this population.

Summary

The study found a significant negative correlation between obesity levels (BMI) and cardiopulmonary capacity (VO₂max) in obese school children aged 8-16 years in Navsari city. Higher BMI values were associated with lower VO₂max, indicating reduced cardiorespiratory fitness in obese children.

Conclusion

In conclusion, the study on the correlation between obesity and cardiopulmonary capacity in school-going children of age 8 to 16 years of Navsari city, India; reinforced strong negative correlation between BMI and maximum oxygen uptake (VO₂max), indicating that higher levels of obesity were associated with reduced cardiopulmonary capacity.

References

- McArdle WD, Katch FI, Katch VL. Exercise physiology: Nutrition, energy, and human performance. Lippincott Williams & Wilkins; c2010.
- Ahirwar R, Mondal PR. Prevalence of obesity in India: A systematic review. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2019 Jan 1;13(1):318-321.
- Freedman DS, Sherry B. The validity of BMI as an indicator of body fatness and risk among children. *Pediatrics*. 2009 Sep;124): S23-34.
- Carayanni V, Bogdanis GC, Vlachopapadopoulou E, Koutsouki D, Manios Y, Karachaliou F, Psaltopoulou T, Michalacos S. Predicting VO₂max in Children and Adolescents Aged between 6 and 17 Using Physiological Characteristics and Participation in Sport Activities: A Cross-Sectional Study Comparing Different Regression Models Stratified by Gender. *Children*. 2022 Dec 9;9(12):1935.
- Buttar KK, Saboo N, Kacker S. A review: Maximal oxygen uptake (VO₂max) and its estimation methods. *Ijpes*. 2019;6(6):24-32.
- Grzyb A, Domagalska-Szopa M, Siwiec A, Kwiecień-Czerwieniec I, Szopa A. Cardiopulmonary Capacity in Overweight and Obese Children and Adolescents: A Cross-Sectional Study. *Frontiers in physiology*. 2021 May 13;12:671827.
- Safaei M, Sundararajan EA, Driss M, Boulila W, Shapi'i A. A systematic literature review on obesity: Understanding the causes & consequences of obesity and reviewing various machine learning approaches used to predict obesity. *Computers in biology and medicine*. 2021 Sep 1;136:104754.
- Himes JH. Challenges of accurately measuring and using BMI and other indicators of obesity in children. *Pediatrics*. 2009 Sep;124): s3-22.
- Morinder G, Mattsson E, Sollander C, Marcus C, Larsson UE. Six-minute walk test in obese children and adolescents: Reproducibility and validity. *Physiotherapy Research International*. 2009 Jun;14(2):91-104.
- Li AM, Yin J, Yu CC, Tsang T, So HK, Wong E, *et al.* The six-minute walk test in healthy children: reliability and validity. *European Respiratory Journal*. 2005 Jun 1;25(6):1057-1056.
- Vanhelst J, Fardy PS, Salleron J, Béghin L. The six-minute walk test in obese youth: reproducibility, validity, and prediction equation to assess aerobic power. *Disability and rehabilitation*. 2013 Mar 1;35(6):479-482.
- Dobrowolska A, Domagalska-Szopa M, Siwiec A, Szopa A. Association between Cardiopulmonary Capacity and Body Mass Composition in Children and Adolescents with High Body Weight: A Cross-Sectional Study. *Children*. 2022 Apr 30;9(5):647.
- Eshwar TM, Chudasama RK, Eshwar ST, Thakrar D. Prevalence of obesity and overweight and their comparison by three growth standards among affluent school students aged 8-18 years in Rajkot. *Indian journal of public health*. 2017 Jan 1;61(1):51.
- Shah H, Mali S, Ranga S, Jadhav C, Rukadikar A, Singh AK, *et al.* Effect of body mass index on cardiorespiratory parameters among medical students: a cross-sectional study. *International Journal of Physiology, Pathophysiology and Pharmacology*. 2022;14(1):4.
- Serés L, López-Ayerbe J, Coll R, Rodríguez O, Manresa JM, Marrugat J, *et al.* Cardiopulmonary function and exercise capacity in patients with morbid obesity. *Revista española de cardiología*. 2003 Jun 1;56(6):594-600.
- Parekh A, Parekh M, Vadasmiya D. Prevalence of overweight and obesity in adolescents of urban & rural area of Surat, Gujarat. *National journal of medical research*. 2012 Sep 30;2(03):325-329.
- Himes JH. Challenges of accurately measuring and using BMI and other indicators of obesity in children. *Pediatrics*. 2009 Sep;124:13-22.
- Directors AB. ATS Statement: Guidelines for the six-minute walk test this official statement of the American Thoracic Society. *Am J Respir. Crit. Care Med*. 2002;166:111-117.
- Racette SB, Yu L, DuPont NC, Clark BR. BMI-for-age graphs with severe obesity percentile curves: tools for plotting cross-sectional and longitudinal youth BMI data. *BMC pediatrics*. 2017 Dec;17(1):1-7.
- Enright PL. The six-minute walk test. *Respiratory care*. 2003 Aug 1;48(8):783-785.
- Morinder G, Mattsson E, Sollander C, Marcus C, Larsson UE. Six-minute walk test in obese children and adolescents: Reproducibility and validity. *Physiotherapy Research International*. 2009 Jun;14(2):91-104.
- Racette SB, Yu L, DuPont NC, Clark BR. BMI-for-age graphs with severe obesity percentile curves: tools for plotting cross-sectional and longitudinal youth BMI data. *BMC pediatrics*. 2017 Dec;17(1):1-7.
- Enright PL, Sherrill DL. Reference equations for the six-minute walk in healthy adults. *American journal of respiratory and critical care medicine*. 1998 Nov 1;158(5):1384-1387.